

Discharge Instructions for Laparoscopic Cholecystectomy Patients

Dr Mark Gately & Dr Martin McLeod

Medications

- Please take regular paracetamol for 7 days after surgery. Tablets or liquid, either are acceptable and DO NOT exceed the maximum daily dosage.
- You will also receive some stronger analgesia eg oxycodone or tapentadol. Take these only if needed. DO NOT DRIVE WHILE TAKING THESE STRONGER ANALGESICS.
- Resume any medications you were on before surgery. Your surgeon will let you know if there are any medications to continue to withhold. This may include blood thinners eg aspirin, warfarin, Plavix etc,

Pain

- It is common to get some abdominal and/or shoulder pain after laparoscopic (key-hole) surgery. This should mainly subside within 3 days of surgery.
- **If the pain worsens or is accompanied by shortness of breath/chest pain, please contact Sea Change Weight Loss Clinic or present to your nearest emergency department.**

Wound Care

- You will leave hospital with dressings on your wounds. Please keep these dressings in-tact for 7 days after surgery.
- You can shower as usual with the dressings on, as they should be waterproof.
- If the dressings come away or get soaked through from the shower, simply replace them with a band-aid
- After 7 days, dressings can be removed. You will see a healing wound. The stitches are beneath the skin and will dissolve and do not need to be removed.
- Contact our office if you have any concerns about your wounds ie: redness, pain, swelling or discharge.
- Note that it is normal to have some redness around wound sites as they heal.

Diet

- You can resume a normal diet however we do advise a lower fat diet for 2 months post laparoscopic cholecystectomy.

Activity

- We would expect most patients will be back at work 1 week after surgery.
- You are free to walk immediately after surgery – start slowly and build up. Avoid too much exercise on hot days and maintain hydration.
- No heavy lifting for 2 weeks following surgery.
- You may drive once you are off strong analgesia and can react in an emergency. Most patients will be driving one week post surgery.

In Case of Emergency

- Please present to an Emergency Department or dial **000** if you experience any of the following:
 - Chest Pain
 - Sudden shortness of breath
 - Fast heart rate – more than 120 beats per min
 - High Temperatures over 38°C
 - Severe Nausea and vomiting
 - Increasing abdominal pain
 - Increasing use of pain medications
 - Inability to pass urine
 - Turning yellow (jaundiced)

Please contact us if you have concerns

- During business hours Mon-Fri 9am-5pm, **contact Sea Change on 02 4037 5858.**
- After hours, contact the ward from which you were discharged.